arXiv:1110.6147v1 [math-ph] 27 Oct 2011

AN ANALYTICAL EVALUATION FOR THE INTEGRAL OF TWO SPHERICAL BESSEL FUNCTIONS WITH AN ADDITIONAL EXPONENTIAL AND POLYNOMIAL FACTOR

R. Mehrem[a]

Associate Lecturer
The Open University in the North West
351 Altrincham Road
Sharston, Manchester M22 4UN
United Kingdom

---

[a] Email: rami.mehrem@btopenworld.com.

ABSTRACT

The integrals $\begin{pmatrix} \lambda_1 & \lambda_2 & \lambda_3 \\ 0 & 0 & 0 \end{pmatrix} \int_0^\infty r^{\lambda_3+1}\, e^{-\alpha r}\, j_{\lambda_1}(k_1 r)\, j_{\lambda_2}(k_2 r)\, dr$ and $\begin{pmatrix} \lambda_1 & \lambda_2 & \lambda_3 \\ 0 & 0 & 0 \end{pmatrix} \int_0^\infty r^{\lambda_3+2}\, e^{-\alpha r}\, j_{\lambda_1}(k_1 r)\, j_{\lambda_2}(k_2 r)\, dr$ are evaluated analytically. The result is a finite sum over the associated Legendre function of the second kind.

1. Introduction

There has been recent interest [1] in the integral $\int_0^\infty r^n\, e^{-\alpha r}\, j_{\lambda_1}(k_1 r)\, j_{\lambda_2}(k_2 r)\, dr$, such that n, λ_1, λ_2 are integers, α, k_1, k_2 are positive real numbers and $n + \lambda_1 + \lambda_2 \geq 0$ to ensure convergence. In the aforementioned reference, a novel symbolic integration method is used in conjunction with Mathematica. The integral has also been solved numerically in another recent reference [2]. In this paper, an alternative approach is shown resulting in a simple expression.

2. Evaluation of the Integral

An earlier result [3] showed that an infinite integral over 3 spherical Bessel functions can be written as

$$\begin{pmatrix} \lambda_1 & \lambda_2 & \lambda_3 \\ 0 & 0 & 0 \end{pmatrix} \int\limits_0^\infty r^2\, j_{\lambda_1}(k_1 r)\, j_{\lambda_2}(k_2 r)\, j_{\lambda_3}(k_3 r)\, dr \;=\; \frac{\pi \beta(\Delta)}{4 k_1 k_2 k_3} i^{\lambda_1+\lambda_2-\lambda_3}$$
$$\times\, (2\lambda_3+1)^{1/2} \left(\frac{k_1}{k_3}\right)^{\lambda_3} \sum_{\mathcal{L}=0}^{\lambda_3} \binom{2\lambda_3}{2\mathcal{L}}^{1/2} \left(\frac{k_2}{k_1}\right)^{\mathcal{L}} \sum_l (2l+1) \begin{pmatrix} \lambda_1 & \lambda_3-\mathcal{L} & l \\ 0 & 0 & 0 \end{pmatrix} \qquad (2.1)$$
$$\times \begin{pmatrix} \lambda_2 & \mathcal{L} & l \\ 0 & 0 & 0 \end{pmatrix} \begin{Bmatrix} \lambda_1 & \lambda_2 & \lambda_3 \\ \mathcal{L} & \lambda_3-\mathcal{L} & l \end{Bmatrix} P_l(\Delta),$$

where $\Delta = (k_1^2 + k_2^2 - k_3^2)/2k_1k_2$, $P_l(x)$ is a Legendre polynomial of order l, $\begin{pmatrix} \lambda_1 & \lambda_2 & \lambda_3 \\ 0 & 0 & 0 \end{pmatrix}$ is a 3j symbol and $\begin{Bmatrix} \lambda_1 & \lambda_2 & \lambda_3 \\ \mathcal{L} & \lambda_3-\mathcal{L} & l \end{Bmatrix}$ is a 6j symbol which can be found in any standard angular momentum text [4, 5]. Using the Closure Relation of the spherical Bessel functions

$$\int\limits_0^\infty k^2\, j_L(kr)\, j_L(kr')\, dk \;=\; \frac{\pi}{2r^2}\, \delta(r - r'), \qquad (2.2)$$

one can write

$$\begin{aligned}\int_0^\infty r^{\lambda_3+1}\, e^{-\alpha r}\, j_{\lambda_1}(k_1 r)\, j_{\lambda_2}(k_2 r)\, dr \;=\; \frac{2}{\pi}\int_0^\infty {k_3}^2\, dk_3\, &\left(\int_0^\infty r^{\lambda_3+1}\, e^{-\alpha r}\, j_{\lambda_3}(k_3 r)\, dr\right) \\ \times\left(\int_0^\infty r^2\, j_{\lambda_1}(k_1 r)\, j_{\lambda_2}(k_2 r)\, j_{\lambda_3}(k_3 r)\, dr\right). \end{aligned} \tag{2.3}$$

From reference [6] one obtains

$$\int_0^\infty r^{\lambda_3+1}\, e^{-\alpha r}\, j_{\lambda_3}(k_3 r)\, dr \;=\; \frac{(2k_3)^{\lambda_3}\,\lambda_3!}{(k_3^2+\alpha^2)^{\lambda_3+1}}. \tag{2.4}$$

Hence,

$$\begin{aligned}\begin{pmatrix}\lambda_1 & \lambda_2 & \lambda_3\\ 0 & 0 & 0\end{pmatrix}\int_0^\infty r^{\lambda_3+1}\, e^{-\alpha r}\, j_{\lambda_1}(k_1 r)\, j_{\lambda_2}(k_2 r)\, dr \;=\; \frac{(2k_1)^{\lambda_3-1}\,(\lambda_3!)}{k_2} i^{\lambda_1+\lambda_2-\lambda_3} \\ \times\,(2\lambda_3+1)^{1/2}\sum_{\mathcal{L}=0}^{\lambda_3}\binom{2\lambda_3}{2\mathcal{L}}^{1/2}\left(\frac{k_2}{k_1}\right)^{\mathcal{L}}\sum_l (2l+1)\begin{pmatrix}\lambda_1 & \lambda_3-\mathcal{L} & l\\ 0 & 0 & 0\end{pmatrix} \\ \times\begin{pmatrix}\lambda_2 & \mathcal{L} & l\\ 0 & 0 & 0\end{pmatrix}\begin{Bmatrix}\lambda_1 & \lambda_2 & \lambda_3\\ \mathcal{L} & \lambda_3-\mathcal{L} & l\end{Bmatrix}\int_0^\infty \beta(\Delta)\,\frac{k_3}{(k_3^2+\alpha^2)^{\lambda_3+1}}\,P_l(\Delta) dk_3. \end{aligned} \tag{2.5}$$

Substituting $k_3^2 \,=\, k_1^2+k_2^2-2k_1k_2\Delta$ and changing the variable of integration from k_3 to Δ, where $k_3 dk_3 \,=\, -k_1k_2 d\Delta$, results in

$$\int_0^\infty \beta(\Delta)\,\frac{k_3}{(k_3^2+\alpha^2)^{\lambda_3+1}}\,P_l(\Delta) dk_3 \;=\; \frac{1}{(2k_1k_2)^{\lambda_3}}\,\frac{(1-y^2)^{-\lambda_3/2}}{\lambda_3!}\,Q_l^{\lambda_3}(y), \tag{2.6}$$

where

$$\int_{-1}^{1}\frac{P_L(x)}{(y-x)^\lambda}\,dx \;=\; \frac{2(1-y^2)^{-(\lambda-1)/2}}{(\lambda-1)!}\,Q_L^{\lambda-1}(y), \tag{2.7}$$

where $Q_L^M(y)$ is the associated Legendre function of the second kind, and $y \,\equiv\, \frac{k_1^2+k_2^2+\alpha^2}{2k_1k_2}$.

The result is

$$\begin{pmatrix} \lambda_1 & \lambda_2 & \lambda_3 \\ 0 & 0 & 0 \end{pmatrix} \int\limits_0^\infty r^{\lambda_3+1}\, e^{-\alpha r}\, j_{\lambda_1}(k_1 r)\, j_{\lambda_2}(k_2 r)\, dr \;=\; \frac{(1-y^2)^{-\lambda_3/2}}{2k_1 k_2^{\lambda_3+1}} i^{\lambda_1+\lambda_2-\lambda_3}$$
$$\times\, (2\lambda_3+1)^{1/2} \sum_{\mathcal{L}=0}^{\lambda_3} \binom{2\lambda_3}{2\mathcal{L}}^{1/2} \left(\frac{k_2}{k_1}\right)^{\mathcal{L}} \sum_l (2l+1) \begin{pmatrix} \lambda_1 & \lambda_3-\mathcal{L} & l \\ 0 & 0 & 0 \end{pmatrix} \tag{2.8}$$
$$\times \begin{pmatrix} \lambda_2 & \mathcal{L} & l \\ 0 & 0 & 0 \end{pmatrix} \begin{Bmatrix} \lambda_1 & \lambda_2 & \lambda_3 \\ \mathcal{L} & \lambda_3-\mathcal{L} & l \end{Bmatrix} Q_l^{\lambda_3}(y).$$

A special case of eq. (2.8) is when $\lambda_3 \,=\, 0$ which leads to

$$\int\limits_0^\infty r\, e^{-\alpha r}\, j_L(k_1 r)\, j_L(k_2 r)\, dr \;=\; \frac{1}{2k_1k_2}\, Q_L(\frac{k_1^2+k_2^2+\alpha^2}{2k_1k_2}). \tag{2.9}$$

As $\alpha \to 0$ and $\lambda_3 = 1$, eq. (2.8) has a singularity when $k_1 = k_2$ as noted in reference [7]. In eq. (2.8), this singularity is due to the factor of $(1-y^2)^{-\lambda_3/2}\, Q_l^{\lambda_3}(y)$. Taking the derivative with respect to α on both sides of eq. (2.4) leads to

$$\int\limits_0^\infty r^{\lambda_3+2}\, e^{-\alpha r}\, j_{\lambda_3}(k_3 r)\, dr \;=\; \frac{2\alpha(2k_3)^{\lambda_3}\,(\lambda_3+1)!}{(k_3^2+\alpha^2)^{\lambda_3+2}}. \tag{2.10}$$

Using the same approach as before one arrives at

$$\begin{pmatrix} \lambda_1 & \lambda_2 & \lambda_3 \\ 0 & 0 & 0 \end{pmatrix} \int\limits_0^\infty r^{\lambda_3+2}\, e^{-\alpha r}\, j_{\lambda_1}(k_1 r)\, j_{\lambda_2}(k_2 r)\, dr \;=\; \frac{\alpha(1-y^2)^{-(\lambda_3+1)/2}}{2k_1^2 k_2^{\lambda_3+2}} i^{\lambda_1+\lambda_2-\lambda_3}$$
$$\times\, (2\lambda_3+1)^{1/2} \sum_{\mathcal{L}=0}^{\lambda_3} \binom{2\lambda_3}{2\mathcal{L}}^{1/2} \left(\frac{k_2}{k_1}\right)^{\mathcal{L}} \sum_l (2l+1) \begin{pmatrix} \lambda_1 & \lambda_3-\mathcal{L} & l \\ 0 & 0 & 0 \end{pmatrix}$$
$$\times \begin{pmatrix} \lambda_2 & \mathcal{L} & l \\ 0 & 0 & 0 \end{pmatrix} \begin{Bmatrix} \lambda_1 & \lambda_2 & \lambda_3 \\ \mathcal{L} & \lambda_3-\mathcal{L} & l \end{Bmatrix} Q_l^{\lambda_3+1}(y). \tag{2.11}$$

Comparing eqs. (2.8) and (2.11) one reaches

$$\int_y^\infty (1-y^2)^{-(L+1)/2}\, Q_l^{L+1}(y)dy \;=\; (1-y^2)^{-L/2}\, Q_l^{L}(y). \tag{2.12}$$

3. Conclusions

Using the Closure Relation of the spherical Bessel functions, an analytical result is obtained for the integrals $\begin{pmatrix} \lambda_1 & \lambda_2 & \lambda_3 \\ 0 & 0 & 0 \end{pmatrix} \int_0^\infty r^{\lambda_3+1}\, e^{-\alpha r}\, j_{\lambda_1}(k_1 r)\, j_{\lambda_2}(k_2 r)\, dr$ and $\begin{pmatrix} \lambda_1 & \lambda_2 & \lambda_3 \\ 0 & 0 & 0 \end{pmatrix} \int_0^\infty r^{\lambda_3+2}\, e^{-\alpha r}\, j_{\lambda_1}(k_1 r)\, j_{\lambda_2}(k_2 r)\, dr$. Work is in progress to obtain integrals of the form $\begin{pmatrix} \lambda_1 & \lambda_2 & \lambda_3 \\ 0 & 0 & 0 \end{pmatrix} \int_0^\infty r^{-\lambda_3}\, e^{-\alpha r}\, j_{\lambda_1}(k_1 r)\, j_{\lambda_2}(k_2 r)\, dr$.

References

1. B. Gebremariam, T. Duguet and S.K. Bogner, Comput. Phys. Comm. **181**, 1136 (2010).
2. J. Van Deun and R. Cools, Comput. Phys. Comm. **178**, 578 (2008).
3. R. Mehrem, J.T. Londergan and M. H. Macfarlane, J. Phys. A **24**, 1435 (1991).
4. D.M. Brink and G.R. Satchler, *Angular Momentum* (Oxford University Press, London 1962).
5. A. R. Edmonds, *Angular Momentum in Quantum Mechanics* (Princeton University Press, Princeton, 1957).
6. I.S. Gradshteyn and I.M. Ryzhik: *Table of Integrals, Series and Products* (Academic Press, New York, 1965).
7. L.C. Maximon, J. Math. Phys. **32**, 642 (1991).